arXiv:1701.00331v1 [math.GR] 2 Jan 2017

ON THE DENSITY OF IMAGES OF THE POWER MAPS IN LIE GROUPS

SAURAV BHAUMIK AND ARUNAVA MANDAL

Abstract. Let G be a connected Lie group. In this paper, we study the density of the images of individual power maps $P_k : G \to G : g \mapsto g^k$. We give criteria for the density of $P_k(G)$ in terms of regular elements, as well as Cartan subgroups. In fact, we prove that if $\mathrm{Reg}(G)$ is the set of regular elements of G, then $P_k(G) \cap \mathrm{Reg}(G)$ is closed in $\mathrm{Reg}(G)$. On the other hand, the weak exponentiality of G turns out to be equivalent to the density of all the power maps P_k. In linear Lie groups, weak exponentiality reduces to the density of $P_2(G)$. We also prove that the density of the image of P_k for G implies the same for any connected full rank subgroup.

1. Introduction

Let G be a connected Lie group and let $\mathfrak{g}$ be the Lie algebra associated to G. The question of whether the exponential map $\exp : \mathfrak{g} \to G$ is surjective or has dense image has been addressed by many authors (see [H], [HM], etc). A connected Lie group G is called *weakly exponential* (resp. *exponential*) if the image of the exponential map exp is dense (resp. surjective). McCrudden [Mc] showed that $\exp : \mathfrak{g} \to G$ is surjective if and only if the *power map* $P_k : G \to G$, defined by $g \to g^k$ is surjective for all $k \geq 2$. The surjectivity of the power maps has been studied by P. Chatterjee ([C1], [C2], [C3], and see there references), R. Steinberg [St], Dani and Mandal [DM]. If a connected Lie group G is weakly exponential, then $P_k(G)$ is dense for all $k \geq 1$.

For an algebraic group over $\mathbb{R}$ or $\mathbb{C}$, the power maps can have dense images without being surjective. For a connected complex algebraic group which is not exponential (e.g., $\mathrm{SL}(2, \mathbb{C})$), there is some $k > 1$ for which P_k is not surjective, but have dense images. One can take Weil restriction to produce such an example over $\mathbb{R}$. For an algebraic group G over $\mathbb{Q}_p$, the question of dense images of the power map is equivalent to the study of surjectivity (see Remark 3.1). The surjectivity of P_k in $G(\mathbb{Q}_p)$ ($\mathbb{Q}_p$-points of the algebraic group G defined over $\mathbb{Q}_p$) has been settled by P. Chatterjee in [C2]. However there does not seem to be any study in the literature on the density of the individual power maps.

This motivated us to consider the following natural question. Given $k > 1$, when is the image $P_k(G)$ dense in G?

On the other hand, weak exponentiality of a Lie group is closely related to the regular elements and Cartan subgroups. A. Borel showed that a connected semisimple Lie group is weakly exponential if and only if all of its Cartan subgroups are connected (see [HM, Theorem 2.10]). Later, K.H. Hofmann showed that for a connected Lie group G (not necessarily semisimple), the set of its regular elements $\mathrm{Reg}(G)$ has a property that, $\mathrm{Reg}(G) \cap \exp(\mathfrak{g})$ is closed in $\mathrm{Reg}(G)$ (see [H, Theorem 17]). Also, he deduced criteria for weak exponentiality of G in terms of regular elements and Cartan subgroups (see [H, Corollary 18]). Following this, K. H. Neeb proved that, a connected Lie group is weakly exponential if and only if all of its Cartan subgroups are connected (see [N, Theorem I.2]).

It is therefore natural to ask whether $P_k(G) \cap \mathrm{Reg}(G)$ is closed in $\mathrm{Reg}(G)$ for all $k \geq 1$.

Key words and phrases. Power maps of Lie groups, regular elements, Cartan subgroups, weak exponentiality, full rank subgroups.

In the following we answer these questions.

Theorem 1.1. *Let G be a connected Lie group. Let $k > 1$ be an integer. The following are equivalent:*

(a) *$P_k(G)$ is dense in G.*
(b) $\mathrm{Reg}(G) \subset P_k(G)$.
(c) *If C is a Cartan subgroup of G, then $P_k(C) = C$.*

Theorem 1.2. *Let G be a connected Lie group. Let $k > 1$ be an integer. Then $P_k(G) \cap \mathrm{Reg}(G)$ is closed in $\mathrm{Reg}(G)$.*

The proof of Theorem 1.1 relies crucially on Theorem 1.2. A result of Wüstner, that every Cartan subgroup is compatible with some Levi decomposition, also turned out to be very useful in many of our proofs.

Analogous results due to Hofmann ([H, Theorem 17, Corollary 18]), and K. H. Neeb ([N, Theorem I.2]) in the context of exponential maps, can be deduced from Theorem 1.2 and Theorem 1.1 (see Corollary 3.2).

In analogy to McCrudden's criterion for exponential images for a connected Lie group, in terms of divisibility, we have the following corollary.

Corollary 1.3. *Let G be a connected Lie group. Then $P_k : G \to G$ is dense for all k if and only if G is weakly exponential.*

The next result also gives a characterization of the density of P_k.

Corollary 1.4. *Let G be a connected Lie group and let C be a Cartan subgroup of G. Let C^* denote the connected component of C and $k > 1$ be an integer. Then $P_k : G \to G$ is dense if and only if $P_k : C/C^* \to C/C^*$ is surjective. In particular, G is weakly exponential if and only if C/C^* is divisible for any Cartan subgroup C.*

The density of k-th power map P_k only depends on its Levi part of the group G. More precisely, if R is the solvable radical (i.e., maximal solvable connected normal subgroup) of G, then $P_k(G)$ is dense in G if and only if $P_k(G/R)$ is dense in G/R (Proposition 3.3). We list some conditions for the density of the images of power maps P_k for simple Lie groups in §4.

The following provides a criterion for weak exponentiality in terms of P_2 only.

Corollary 1.5. *Let G be a connected linear Lie group. Then G is weakly exponential if and only if $P_2 : G \to G$ is dense.*

This means that (compare with Theorem 1.1), in a connected linear Lie group G, if every element that belongs to a Cartan subgroup admits a square root then G is weakly exponential. For linear groups, this extends Neeb's result, which says that G is weakly exponential if and only if all of its Cartan subgroups are connected. On the other hand it strengthens one of P. Chatterjee's results ([C3, Theorem 1.6]). However, the corollary is no longer valid for non-linearizable groups, as we observe in Example 5.1.

The following can be thought of as a power map analogue of Neeb's result (see [N, Proposition I.6]).

Theorem 1.6. *Let G be a connected Lie group and let A be a connected full rank subgroup of G. Let $k > 1$ be an integer. Then $P_k : G \to G$ is dense implies $P_k : A \to A$ is dense.*

Let G be a connected linear reductive Lie group and $\mathfrak{g}$ be its associated Lie algebra. Let θ be the Cartan involution on $\mathfrak{g}$. Then there is a natural Cartan decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$, where $\mathfrak{k}$ and $\mathfrak{p}$ are the eigenspaces corresponding to the eigenvalues 1 and -1 respectively.

It follows that $[\mathfrak{k},\mathfrak{k}] \subseteq \mathfrak{k}$, $[\mathfrak{k},\mathfrak{p}] \subseteq \mathfrak{p}$, $[\mathfrak{p},\mathfrak{p}] \subseteq \mathfrak{k}$ and $\mathfrak{z}(\mathfrak{g}) \subseteq \mathfrak{k}$. Let $\mathfrak{a}$ be a maximal abelian subspace of $\mathfrak{p}$ and take a Cartan subalgebra t_a of $\mathfrak{z}_{\mathfrak{k}}(\mathfrak{a})$. Define $\mathfrak{c} := t_a + \mathfrak{a}$. It is noted in Neeb [N] that $\mathfrak{c}$ is a Cartan subalgebra of $\mathfrak{g}$ and every Cartan subalgebra $\mathfrak{c}'$ of $\mathfrak{g}$ is conjugate to a Cartan subalgebra is of the form $t'_a + \mathfrak{a}'$, where $\mathfrak{a}' \subseteq \mathfrak{a}$ and $t'_a \supseteq t_a$.

We give below another criterion for the density of images of power maps in case of reductive groups, which is mostly important for non-split case.

Corollary 1.7. *Let G be a connected reductive Lie group. Let $k > 1$ be an integer. Then $P_k : G \to G$ is dense if and only if $P_k : Z_G(t_a) \to Z_G(t_a)$ is dense.*

The paper is organized as follows. We recall some definitions and prove some results on regular elements, Cartan subgroups, including Theorem 1.2, in §2. Theorem 1.1, Corollaries 1.3, 1.4 are proved in §3. We discuss the density of power maps in simple Lie groups in §4. Corollary 1.5 is proved in §5. Finally, in §6 we prove Theorem 1.6 and Corollary 1.7.

2. Regular elements, Cartan subgroups and power maps

In this section we recall some definitions, introduce notations, prove a few lemmas and Theorem 1.2.

Definition 1. [H] An element g in a Lie group G is called *regular* if the nilspace $N(\mathrm{Ad}_g - 1) \subset \mathfrak{g}$ is of minimal possible dimension. The set of regular elements of G is denoted by Reg(G).

Definition 2. [C1] Let G be a Lie group. An element g in G is called *P_k-regular* if dP_k is nonsingular at g.

Lemma 2.1. *Let G be a real Lie group. Suppose $g, h \in G$, and $h^k = g$. Then g is regular if and only if h is both regular and P_k-regular.*

Proof. Let g be regular. Writing $T = \mathrm{Ad}_g - 1$, $U = \mathrm{Ad}_h - 1$ and

$$V = \mathrm{Ad}_{h^{(k-1)}} + \mathrm{Ad}_{h^{(k-2)}} + \ldots + \mathrm{Ad}_h + 1,$$

we get $T = UV$. Then $N(U) \subset N(T)$, and by minimality of $\dim(N(T))$, we conclude that h is regular.

Now, the complexification $\mathfrak{g}_{\mathbb{C}}$ of the real Lie algebra $\mathfrak{g}$, splits into direct sum of generalized eigenspaces V_α under Ad_h, i.e., $\mathfrak{g}_{\mathbb{C}} = \oplus_{\alpha\in\Delta} V_\alpha$ (where Δ denotes the set of eigenvalues of Ad_h). We note that $V_1 = N(U) \otimes \mathbb{C}$. Let $\alpha^k = 1$ but $\alpha \neq 1$. Then in the splitting of $\mathfrak{g}_c$ under $\mathrm{Ad}_h^k = \mathrm{Ad}_g$, the generalized eigenspace for 1 will contain $V_1 \oplus V_\alpha$. By minimality of $\dim_{\mathbb{C}}(V_1) = \dim_{\mathbb{R}}(N(U))$, $V_\alpha = 0$. By [C1, Lemma 2.1], it follows that h is P_k-regular.

Conversely, let h be P_k-regular and regular. By [C1, Lemma 2.1], V is invertible on $\mathfrak{g}_{\mathbb{C}}$ and $N(U) \otimes \mathbb{C} = N(T) \otimes \mathbb{C}$ and hence $N(T)$ is of minimal dimension. □

We now recall the definition of Cartan subgroup of a Lie group. Let G be a connected Lie group with its Lie algebra $\mathfrak{g}$. Let $\mathfrak{c}$ be the Cartan subalgebra of $\mathfrak{g}$ and let Δ be the set of roots of $\mathfrak{g}_{\mathbb{C}}$ belonging to $\mathfrak{c}_{\mathbb{C}}$. Thus $\mathfrak{g}_{\mathbb{C}} = \mathfrak{c}_{\mathbb{C}} + \Sigma_{\alpha\in\Delta}\mathfrak{g}^\alpha_{\mathbb{C}}$. The normalizer $N_G(\mathfrak{c})$ of $\mathfrak{c}$ is defined by $N_G(\mathfrak{c}) := \{g \in G | \mathrm{Ad}_g(\mathfrak{c}) = \mathfrak{c}\}$. Note that $N_G(\mathfrak{c})$ is a closed subgroup of G with Lie algebra $\mathfrak{n}_{\mathfrak{g}}(\mathfrak{c}) = \{X \in \mathfrak{g} | [X, \mathfrak{c}] \subseteq \mathfrak{c}\}$. Let $C(\mathfrak{c}) := \{g \in N_G(\mathfrak{c}) | \alpha \circ \mathrm{Ad}_g|_{\mathfrak{c}_{\mathbb{C}}} = \alpha, \forall \alpha \in \Delta\}$.

A closed subgroup C of a Lie group G is called a *Cartan subgroup* if its Lie algebra $\mathfrak{c}$ is a Cartan subalgebra of $\mathfrak{g}$ and $C(\mathfrak{c}) = C$. This agrees with the usual definition of a Cartan subgroup (see Appendix of [N]). In particular, a Cartan subgroup is nilpotent.

Let us recall a theorem due to M. Wüstner ([W, Theorem 1.9 (ii)]). Let G be a connected Lie group and let R be its solvable radical. Then for any Cartan subgroup C of G, there is

a Levi subgroup S of G such that C can be decomposed as $C = (C \cap S)(C \cap R)$. Moreover, $C \cap S$ is a Cartan subgroup of S, $C \cap R$ is connected and $C \cap R \subseteq Z_G(S)$.

Now we can immediately observe that, if G is connected linear Lie group then any Cartan subgroup can be written as $C = TN$ (almost direct product), where T is abelian and N is connected nilpotent subgroup of C. Indeed, if G is linear, so is S. This means, S is isomorphic to the identity component (in Euclidean topology) of a geometrically connected semisimple algebraic group defined over $\mathbb{R}$. In this case, $T = C \cap S$ is abelian, and $N = C \cap R$ is connected nilpotent. This will be used later.

Lemma 2.2. *Let G be a Lie group, and let $Z \subset Z(G)$ be a closed central subgroup. Let $G_1 = G/Z$, and let $\pi : G \to G_1$ be the projection.*

(a) If C is a Cartan subgroup of G, then $C_1 = \pi(C)$ is a Cartan subgroup of G_1, and $C = \pi^{-1}(C_1)$.

(b) Suppose Z is connected. If $P_k(C_1)$ is both closed and open in C_1, then $P_k(C)$ is both closed and open in C.

Proof. (a) It follows from the description of Cartan subgroup in terms of its adjoint action on the Lie algebra. Indeed, if $\mathfrak{c}$ is a Cartan subalgebra of $\mathfrak{g}$, then $\mathfrak{c}_1 = \mathfrak{c}/\mathfrak{z}$ is a Cartan subalgebra of $\mathfrak{g}_1 = \mathfrak{g}/\mathfrak{z}$, while $\mathfrak{z} \subset \mathfrak{z}(\mathfrak{g}) \subset \mathfrak{z}_{\mathfrak{g}}(\mathfrak{c})$. For any $x \in G$, Ad_x preserves $\mathfrak{z}(\mathfrak{g})$, hence $\mathfrak{z}$. Therefore $\mathrm{Ad}_{\pi(x)}$ normalizes $\mathfrak{c}_1$ if and only if Ad_x normalizes $\mathfrak{c}$. The roots $\overline{\alpha}$ of $\mathfrak{g}_1$ with respect to $\mathfrak{c}_1$ are induced from roots α of $\mathfrak{g}$ with respect to $\mathfrak{c}$, and $\mathfrak{g}^{\alpha} = \mathfrak{g}_1^{\overline{\alpha}}$. Hence C_1 is a Cartan subgroup of G_1.

Again, $C \subset \pi^{-1}(C_1)$. Let $x \in \pi^{-1}(C_1)$. Since $C_1 = \pi(C)$, there is some $y \in C$ such that $\pi(x) = \pi(y)$. This means, $x = yz$, for some $z \in Z \subset Z(G) \subset C$, which proves $C = \pi^{-1}(C_1)$.

(b) By (a), we get a short exact sequence $1 \to Z \to C \xrightarrow{\pi} C_1 \to 1$. Now, since Z is a connected abelian group, for any $x \in C$, $xZ \cap P_k(C)$ is nonempty if and only if $xZ \subset P_k(C)$. Therefore $P_k(C) = \pi^{-1}(P_k(C_1))$. □

Lemma 2.3. *Let C be a Lie group and let $\tilde{C}$ be a cover of C. If $P_k(\tilde{C})$ is both open and closed then so is $P_k(C)$.*

Proof. Let $\pi : \tilde{C} \to C$ be the covering map. Then $\pi(P_k(\tilde{C})) = P_k(C)$. By hypothesis, $P_k(\tilde{C})$ is disjoint union of connected components of $\tilde{C}$. Let $P_k(\tilde{C}) = \sqcup_{i \in I} \tilde{X}_i$, where $\tilde{X}_i$'s are connected component of $\tilde{C}$ and I is the indexing set. For any i, $\pi(\tilde{X}_i)$ is connected, hence there is a connected component X_i of C such that $\pi(\tilde{X}_i) \subseteq X_i$. We claim that for $i \in I$, $\pi(\tilde{X}_i)$ is a connected component of C. Suppose the claim holds, then $\pi(\tilde{X}_i) = X_i$ and $\pi(P_k(\tilde{C})) = \sqcup_{i \in I} \pi(\tilde{X}) = \sqcup_{i \in I} X_i$. As X_i's are both open and closed in C, $P_k(C)$ is both open and closed. This will prove the lemma.

To prove our claim, we note that, as π is a covering map, $\pi(\tilde{X}_i)$ is open in X_i, therefore it is enough to prove that $\pi(\tilde{X}_i)$ is closed in X_i. Now, let x be an element in the closure of $\pi(\tilde{X}_i)$. As π is covering of a Lie group C, there is a *connected* open set U containing x such that $\pi^{-1}(U) = \sqcup_r V_r$, where V_r's are open in $\tilde{X}$ and π restricted to V_r is a diffeomorphism onto U. Then, there exists r_0, such that $\tilde{X}_i \cap V_{r_0} \neq \emptyset$. Since V_{r_0} is connected, it must lie entirely in the connected component $\tilde{X}_i$. Therefore $x \in U = \pi(V_{r_0}) \subset \pi(\tilde{X}_i)$. □

Proposition 2.4. *Let G be any connected real Lie group and let C be a Cartan subgroup of G. For any $k \geq 0$, $P_k(C)$ is both open and closed in C.*

Proof. First note that it is enough to consider the case when $Z(G)$ is discrete. Indeed, let $G_1 = G/Z(G)^*$, where $Z(G)^*$ is the identity component of $Z(G)$, and let $\pi_1 : G \to G_1$ be the projection. Then by Lemma 2.2, it is enough to prove that $P_k(C_1)$ is closed and open in C_1. Inductively, $G_{i+1} = G_i/Z(G_i)^*$, and let C_{i+1} be the image of C_i in G_{i+1}. Since the

dimension of G is finite, there is some i such that $Z(G_i)$ is discrete. Henceforth, we will assume that $Z = Z(G)$ is discrete.

Note that G is a covering group of the linear group $G' = G/Z$. Let $\pi : G \to G'$ denote the covering projection. Therefore, by Lemma 2.2, $\pi(C)$ is a Cartan subgroup of G'. Then by Wüstner, there is a Levi subgroup S of G' such that $\pi(C) = (\pi(C) \cap S)(\pi(C) \cap R)$ (almost direct product), where R is the solvable radical of G'.

Now, write $G' = S \cdot R$ (almost semidirect product). As S acts on R, we consider an external semidirect product of S and R, denoted by $S \ltimes R$. We observe that there is a surjective group homomorphism $P : S \ltimes R \to S \cdot R$, whose kernel is isomorphic to $D = \{(z, z^{-1}) | z \in S \cap R\}$. Since D is discrete normal, it is central in $S \ltimes R$. Hence by Lemma 2.2, $P^{-1}(\pi(C))$ is a Cartan subgroup of $S \ltimes R$ and in fact $P^{-1}(\pi(C)) = (\pi(C) \cap S) \times (\pi(C) \cap R)$. As $\pi(C) \cap R$ is connected linear nilpotent, it can be written as $T_1 \times N$ for some torus T_1 and simply connected subgroup N of $\pi(C) \cap R$. Therefore $P^{-1}(\pi(C))$ is of the form $T \times N$, where $T = (\pi(C) \cap S) \times T_1$. Hence $P_k(P^{-1}(\pi(C))$ is both open and closed in $P^{-1}(\pi(C))$ implies $P_k(\pi(C))$ is so in $\pi(C)$, as $S \ltimes R$ is a finite covering of G'. Therefore, to prove $P_k(\pi(C))$ is both open and closed, it is enough to show the same for $P_k(P^{-1}(\pi(C)))$.

Hence, by Lemma 2.3, we may assume that $\pi(C) = T \times N$, where T is an abelian group and N is a simply connected nilpotent group. Now, $\pi : C \to T \times N$ is a covering projection. Therefore C has to be of the form $A \times N$, where A is a covering space of T. Indeed, let $H_1 = \pi^{-1}(T \times \{1\})$ and $H_2 = \pi^{-1}(\{1\} \times N)$, where $\{1\}$ denotes the identity element in G. Then H_1, H_2 are normal in C. Again, N is simply connected, so the covering space $H_2 \to N$ has a section, while the kernel Z is central. Hence $H_2 = N \times Z$. Now, let $x \in H_1$. Then ι_x induces an action on H_2 which preserves the central Z. Let $(y, z) \in N \times Z$. Then $\iota_x(y, z) = (y', z')$. Now, taking projection to N, since $x \in H_1$, we see that $y = y'$. Hence the action of ι_x is given by a matrix

$$\begin{pmatrix} id_N & \phi \\ \{1\} & id_Z \end{pmatrix}$$

where ϕ is a homomorphism $N \to Z$. Now, since N is connected and Z is discrete, ϕ is zero, hence H_1 and H_2 commute. Therefore we have a surjective homomorphism of groups $H_1 \times H_2 \to C$ given by inclusions and multiplication in C. The kernel is $\{(z, z^{-1}) | z \in Z\}$. Since $H_2 = N \times Z$, we see that the induced homomorphism $H_1 \times N \to C$ is an isomorphism. We set, $A = H_1$.

This means, it is enough to prove that $P_k(A)$ is closed in A. So we can assume that C is a covering group of T with central kernel Z.

Since G' is linear, T is of the form $F \times T_2$, where F is a finite $\mathbb{Z}/2\mathbb{Z}$-vector space, and T_2 is a connected abelian group. Now, in order to prove that $P_k(C)$ is both open and closed in C, by Lemma 2.3, it is enough to do so after replacing C by a covering group. In particular, we can replace T_2 by its simply connected covering, which will be a vector space, say V, and replace C by the fiber product of V and C over T_2.

So we can assume that C is an extension of $F \times V$ by the central Z.

Let $p : C \to F$ be the composite $C \xrightarrow{\pi} F \times V \xrightarrow{Pr_1} F$. Let us consider the normal subgroup $B = \pi^{-1}(0 \times V)$. This is a covering group of the simply connected V by Z, hence is of the form $Z \times V$. Since V is connected abelian and Z is discrete, B is central by similar argument as above.

Therefore, we can express C as an extension of F by the central $B = Z \times V$ as follows.

$$0 \to B \to C \xrightarrow{p} F \to 0.$$

Let the number of elements in F be r, and let $F = \{p(x_1), \ldots, p(x_r)\}$. Now, C is the disjoint union of the cosets $x_i B$. Therefore, $P_k(C)$ is the union of $P_k(x_i B)$. Note that,

since B is central in C, $P_k(xB) = x^k P_k(B)$, and hence is closed and open in $x^k B$. This implies that $P_k(C) = \cup_{i=1}^{r} P_k(x_i B)$ is closed and open in C. □

Proof of Theorem 1.2: By [B, Ch. VII, §4 n° 2, Proposition 5], for each regular $g \in G$ and for each identity neighbourhood W there is a neighbourhood V of g such that for each $x \in V$ there is a $w \in W$ with $C(x) = \iota_w C(g)$ (equivalently, $\mathfrak{c}(x) = \mathrm{Ad}_w \mathfrak{c}(g)$ at the level of Lie subalgebras).

Let $g \in \mathrm{Reg}(G)$ and let $g_n \in \mathrm{Reg}(G)$ be a sequence such that $g_n = h_n^k$ and g is the limit of the sequence i.e. $g_n \to g$ in G. Note that $P_k : G \to G$ is nonsingular at identity, because dP_k at identity is multiplication by k. Hence there are sequences of identity neighbourhoods W_i, W_i' such that both the diameters of W_i, W_i' tend to zero, and $W_i = P_k(W_i')$. Now, let V_i be neighbourhoods of g corresponding to W_i'. Since $g_n \to g$, for each i there is some $g_{n_i} \in V_i$. We can choose $n_{i+1} > n_i$. Hence we can assume without loss of generality that $g_i \in V_i$. We can therefore find $\eta_i \in W_i'$ such that ${\eta_i}^{-1} C(g_i) \eta_i = C(g)$. Since $C(g_i) = C(h_i)$, ${\eta_i}^{-1} h_i \eta_i \in C(g)$ for all i, and hence writing $\xi_i = \eta_i^k \in W_i$, ${\xi_i}^{-1} g_i \xi_i \in P_k(C(g))$. Since $\xi_i \to 1$ and $g_i \to g$, ${\xi_i}^{-1} g_i \xi_i \to g$. Hence, by closedness of $P_k(C(g))$ (Proposition 2.4), we get $g \in P_k(C(g))$. □

3. Characterization of the density of the power map P_k

In this section we prove Theorem 1.1 by using Theorem 1.2. Also, we deduce many characterization about the dense images of the power map, which can be used in the rest of the paper.

Proof of Theorem 1.1: (a) ⇔ (b) If $P_k(G)$ is dense in G then $P_k(G) \cap \mathrm{Reg}(G)$ is dense in $\mathrm{Reg}(G)$, hence equal to $\mathrm{Reg}(G)$, by Theorem 1.2. As $\mathrm{Reg}(G)$ is dense in G and it is contained in $P_k(G)$, the converse follows.

(b) ⇔ (c) Suppose $\mathrm{Reg}(G) \subset P_k(G)$. It is enough to show that $C \cap \mathrm{Reg}(G) \subset P_k(C)$, because $C \cap \mathrm{Reg}(G)$ is dense in C (see [H1, Proposition 1.6]), while $P_k(C)$ is closed in C (Proposition 2.4). If $x \in C \cap \mathrm{Reg}(G)$ then by assumption, there is $y \in G$ such that $x = y^k$. This implies $y \in \mathrm{Reg}(G)$, by Lemma 2.1. Then y lies in a unique Cartan subgroup, which has to be C because $y^k \in C$. Conversely, any regular element g can be found in a unique Cartan subgroup C. □

Remark 3.1. For algebraic groups we can give a more straightforward argument.

(1) Let G be an algebraic group over a field $\mathbb{F}$. Let C be a Cartan subgroup of G. Then for $\mathbb{F} = \mathbb{R}$ or $\mathbb{Q}_p$, the regular elements of $G(\mathbb{F})$ (group of $\mathbb{F}$-rational point of the algebraic group G) which can be conjugated to $C(\mathbb{F})$ form an open subset of $G(\mathbb{F})$ in the Euclidean or the ultrametric topology.

To see this, let $\Phi : G \times C \to G : (h, y) \mapsto hyh^{-1}$ and fixing $y \in C$, let $\Phi_y : G \to G$ is given by $\Phi_y(h) = hyh^{-1}$. Note that the differential of Φ at (e, y), where $y \in \mathrm{Reg}(G) \cap C$, is surjective. Write $T_y C = dL_y T_e C$,

$$d\Phi(\xi, dL_y v) = d\Phi(\xi, 0) + dL_y v = d\Phi_y(\xi) + dL_y v = dL_y(df(\xi) + v),$$

where $f : G \to G : h \mapsto y^{-1} hyh^{-1}$. Note that, f is the composition of $G \to G \times G : h \mapsto (y^{-1}hy, h^{-1})$ and $G \times G \overset{\mu}{\to} G : (x, y) \mapsto xy$. Then,

$$df(\xi) = d\mu(\mathrm{Ad}_y{}^{-1}\xi, -\xi) = (\mathrm{Ad}_y{}^{-1} - I)\xi.$$

Now, since $y \in C$ is regular, $T_e C = N(\mathrm{Ad}_y{}^{-1} - I)$. Therefore we get the following.

$$d\Phi(\mathfrak{g} \times T_y C) = dL_y(\mathrm{im}(\mathrm{Ad}_y{}^{-1} - I) + N(\mathrm{Ad}_y{}^{-1} - I)) = \mathfrak{g}.$$

(2) Now, for $\mathbb{F} = \mathbb{Q}_p$, if the group is not anisotropic, there is a nonempty open subset of regular elements which can be conjugated to a Cartan subgroup C which contains a copy of $\mathbb{Q}_p^\times = \mathbb{Z} \times \mathbb{Z}_p^\times$. In this case, for any $k > 0$, $P_k(C)$ is proper, open and closed

in C. Therefore none of (a), (b), (c) of Theorem 1.1 will hold. On the other hand, if G is anisotropic then it is weakly exponential. In this case, density of P_k is equivalent to surjectivity.

(3) For $\mathbb{F} = \mathbb{R}$, if the group contains any disconnected Cartan subgroup C, then $P_{2k}(C) = C^*$, the identity component. Therefore, P_{2k} is never dense. On the other hand, if $P_{2k+1}(C) = C$, so that $P_{2k+1}(G)$ contains all regular elements, hence is dense. In case all Cartan subgroups are connected G is weakly exponential.

Proof of Corollary 1.3: Suppose that $P_k(G)$ is dense in G. Then by Theorem 1.1, $P_k(G)$ contains a dense open set (namely $\mathrm{Reg}(G)$). Hence by Baire category theorem, $\cap_{k\geq 1}P_k(G)$ contains a non empty open set and therefore G is weakly exponential. Now the converse follows from McCrudden criterion. □

Theorem 1.1 can be used to obtain the following criteria of weak exponentiality, which is well known ([N, Theorem I.2]).

Corollary 3.2. *Let G be a connected Lie group and let $\mathfrak{g}$ be the Lie algebra associated to G. The following are equivalent.*

(a) *G is weakly exponential.*
(b) $\mathrm{Reg}(G) \subseteq \exp(\mathfrak{g})$.
(c) *All Cartan subgroups of G are connected.*

Proof. $(a) \Leftrightarrow (b)$ By Corollary 1.1, G is weakly exponential if and only if $P_k(G)$ is dense for all k, which is equivalent to saying that $\mathrm{Reg}(G) \subseteq \cap_{k\geq 1}P_k(G) \subseteq \exp(\mathfrak{g})$, by Theorem 1.1.

$(a) \Leftrightarrow (c)$ By using Theorem 1.1, G is weakly exponential if and only if all Cartan subgroups of G are divisible (i.e., $P_k(C) = C$ for all $k \geq 1$). By [HL, Proposition 1], this is equivalent to saying that C is connected. □

Proposition 3.3. *Let G be a connected Lie group, and let R be the radical of G. Let $k > 0$ be an integer. Then $P_k : G \to G$ is dense if and only if $P_k : G/R \to G/R$ is dense.*

Proof. The right implication is obvious. For the converse, let H be a Cartan subgroup of G. Then there exists a Levi subgroup S of G such that $H = (H \cap S)(H \cap R)$. We write, $G = S \cdot R$. Let $Z = S \cap R$ and $\Delta(Z) := \{(z, z^{-1}) | z \in Z\}$. Now, consider $S \ltimes R$ as an external semidirect product of S and R. The map $\pi : S \ltimes R \to S \cdot R$ sending $(s, r) \to s \cdot r$, for $s \in S$, $r \in R$, is a covering map and G can be thought of as a quotient of $S \ltimes R$ by $\Delta(Z)$. Then Cartan subgroup $\pi^{-1}(H)$ of $S \ltimes R$ is of the form $C \times N$ where $C = H \cap S$ is a Cartan subgroup of S, $N = H \cap R$ is connected nilpotent subgroup of R, and $H = (C \times N)/\Delta(Z)$. Also, note that as $S \to S/Z$ is covering, then C/Z is a Cartan subgroup of G/R. Now, there is a short exact sequence of groups

$$1 \to N \to (C \times N)/\Delta(Z) \to C/Z \to 1.$$

As $P_k(G/R)$ is dense in G/R, by Theorem 1.1, $P_k(C/Z) = C/Z$. Now, let $x = \overline{x_1 x_2} \in (C \times N)/\Delta(Z)$ for some $x_1 \in C$ and $x_2 \in N$. As $P_k(C/Z) = C/Z$, there exists $y_1 \in C$ and $z \in Z$ such that $x_1 = y_1^k z$. Thus $\overline{x_1 x_2} = \overline{y_1^k z x_2} = y_1^k y_2^k$ in H, for some $y_2 \in N$, as z is contained in N and N is connected nilpotent Lie group. Now we notice that $\overline{x_1 x_2} = \overline{y_1 y_2}^k$, as C and N commute. Hence the power map $P_k : H \to H$ is surjective. Therefore, by applying Theorem 1.1, the proposition follows. □

To deduce Corollary 1.4, we need next two lemmas.

Lemma 3.4. *Let $\pi : G \to H$ be a covering of connected Lie groups, with discrete kernel D. Let $C \subset H$ be a connected Cartan subgroup which is abelian. Then the corresponding Cartan subgroup $M = \pi^{-1}(C)$ is also abelian.*

Proof. Note that the identity component M^* is a covering of C. Since C is connected abelian, M^* is abelian. Any element $x \in M$ is of the form $x = mz$ for some $m \in M^*$ and $z \in D$. Now we note that, the group homomorphism $M^* \times D \to M$, given by $(m', z') \to m'z'$, is surjective. Hence M is abelian. □

Lemma 3.5. *Let $\tilde{G}$ be a connected Lie group with discrete center. Let $\tilde{C}$ be any Cartan subgroup of $\tilde{G}$. Then the connected component $\tilde{C}^*$ of $\tilde{C}$ is central.*

Proof. Let $G = \tilde{G}/Z(\tilde{G})$ and $C := \pi(\tilde{C})$ be the corresponding Cartan subgroup of G. We can write $C = A \times F$, where A is connected and F is a $\mathbb{Z}/2\mathbb{Z}$ vector space. If F is trivial, by Lemma 3.4, $\tilde{C}$ is abelian and hence the conclusion holds. Suppose that F is not trivial. We observe that $\pi^{-1}(F)$ is a normal subgroup of $\tilde{C}$, where $\pi : \tilde{C} \to C$ is the natural projection map. Therefore, $\tilde{C}^*$ acts trivially on $\pi^{-1}(F)$, as $\tilde{C}^*$ is connected and $\pi^{-1}(F)$ is discrete. Thus elements of $\tilde{C}^*$ and $\pi^{-1}(F)$ commute each other. Note that $\tilde{C}$ is a quotient of $\tilde{C}^*$ and $\pi^{-1}(F)$. Indeed, for $x \in \tilde{C}$, let $\pi(x) = (a, b) \in A \times F$. We can choose $x_1 \in \tilde{C}^*$ such that $x_1 = \pi^{-1}(a)$. Then it follows that $xx_1^{-1} \in \pi^{-1}(F)$. Thus $x = (xx_1^{-1})x_1 \in \pi^{-1}(F) \times \tilde{C}^*$ and hence $\tilde{C}^*$ is central in $\tilde{C}$. □

Proof of Corollary 1.4: ($:\Rightarrow$) This follows from Theorem 1.1.

($\Leftarrow:$) Let R be the solvable radical of G. We recall that, there is a decomposition of the Cartan subgroup C ([W, Theorem 1.9 (ii)]). For C, there is a Levi subgroup S of G such that $C = C_s \cdot N$ (almost direct product), where $C_s = C \cap S$ is a Cartan subgroup of S and $N = C \cap R$ is connected nilpotent contained in $Z_G(S)$. Therefore we observe that $C/C^* \simeq C_s/C_s^*Z$, where $Z = C_s \cap N$, a central subgroup. Also, we note that if C_1 is a Cartan subgroup of G/R, then $C_1/C_1^* \simeq C_s/C_s^*Z$. Hence $P_k : C/C^* \to C/C^*$ is surjective if and only if $P_k : C_1/C_1^* \to C_1/C_1^*$ is surjective. Applying Lemma 3.5 for the semisimple group G/R, we get $P_k : C_1 \to C_1$ is surjective and hence by Theorem 1.1, $P_k : G/R \to G/R$ is dense. Now by Proposition 3.3, we conclude that $P_k(G)$ is dense in G. □

4. Density of the power maps on simple Lie groups

In this section we study the dense images of the power maps in simple Lie groups. We use results in §3, to determine the integer k, for which P_k has dense images.

We recall from Remark 3.1 that, for a connected real algebraic group G with disconnected Cartan subgroup, P_k is dense if and only if k is odd. However, in general this is not true.

Example 4.1. Let $\tilde{G} = \widetilde{\mathrm{SL}(2,\mathbb{R})}$ be the simply connected cover of $\mathrm{SL}(2,\mathbb{R})$. Then $P_k : \widetilde{\mathrm{SL}(2,\mathbb{R})} \to \widetilde{\mathrm{SL}(2,\mathbb{R})}$ is not dense for all $k > 1$. Indeed, the center of $\tilde{G}$ is infinite cyclic, and the adjoint group $\mathrm{PSL}(2,\mathbb{R})$ has a Cartan subgroup isomorphic to $\mathbb{R}$. The corresponding Cartan subgroup in $\tilde{G}$ is therefore isomorphic to $\mathbb{R} \times \mathbb{Z}$, where the component $\mathbb{Z}$ is contributed by the center. Therefore for $k > 1$, P_k is never surjective on this Cartan subgroup, so the assertion follows by Theorem 1.1.

There is a list of weakly exponential non-compact non-complex simple Lie algebras in Neeb ([N]). Neeb also mentioned exactly which in the list are completely weakly exponential (i.e., whose corresponding simply connected group is weakly exponential). Thus recalling Borel's criterion about weak exponentiality, one can understand the connectedness of the Cartan subgroup.

For a connected Lie group G, if Cartan subgroups are connected, then P_k is dense in G for all k, as G is weakly exponential. Therefore we concentrate the case when Cartan subgroups are disconnected. As we noted before (Proposition 3.3), the study of density

boils down to semisimple Lie group, at this moment we now investigate this phenomenon for simple Lie groups.

Let us fix some notations for this section. Let $\tilde{G}$ be the universal cover of the connected simple adjoint Lie group G. Let π denote the covering map and $\tilde{C}$ be a Cartan subgroup of $\tilde{G}$. Then let $C = \pi(\tilde{C})$ be the corresponding Cartan subgroup of G. Let us denote the fundamental group of G by $\pi_1(G)$. So we get a natural short exact sequence

$$1 \to \pi_1(G) \to \tilde{C} \to C \to 1.$$

For a connected simple adjoint Lie group G, we notice that its fundamental group is finitely generated, whose possible factors are isomorphic to $\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z}$, $\mathbb{Z}/4\mathbb{Z}$, $\mathbb{Z}/8\mathbb{Z}$, or $\mathbb{Z}/3\mathbb{Z}$. We discuss some cases. In the following cases we assume that some Cartan subgroup of $\tilde{G}$, say $\tilde{C}$ is disconnected.

Case 1. Let $\pi_1(G) \simeq \mathbb{Z}_2$ or non cyclic group of order 4.

This happens for A_nI ($n \geq 2$), B_nI, E_6^6I, E_8^8VIII, F_4^4I, G_2^2I, C_nII, $E_7^{-5}VI$, $E_8^{-24}IX$, $F_4^{-22}II$. Note that none of these is weakly exponential.

Lemma 4.1. *In Case 1, $P_k : \tilde{G} \to \tilde{G}$ is dense if and only if k is odd.*

Proof. First assume that the Cartan subgroup $C = \pi(\tilde{C})$ of G is disconnected. By hypothesis, it is enough to check for odd k, as for even k, $P_k(G)$ is not dense in G. Note that $\pi_1(G)$ is central in $\tilde{C}$. Since for k odd, P_k is surjective on C and $\pi_1(G)$ as well, $P_k : \tilde{C} \to \tilde{C}$ is surjective and hence by Theorem 1.1, the lemma follows.

Now, suppose that C is connected. Using the same argument as before, for odd k, P_k is dense in $\tilde{G}$. Now Suppose that P_2 is dense in $\tilde{G}$. Then P_k is dense in $\tilde{G}$ for all k, which further implies that $P_k(\tilde{C}) = \tilde{C}$ for all k, by Theorem 1.1. Thus by [HL, Proposition 1], $\tilde{C}$ is connected, which is a contradiction. This implies that, for even k, $P_k(\tilde{G})$ is not dense in $\tilde{G}$, as desired. □

Case 2(a). C is connected, and $\pi_1(G) \simeq \mathbb{Z}$. This occurs for $E_7^{-25}VII$, A_1I.

As C is connected, we get the following short exact sequence

$$1 \to \tilde{C}^* \cap \mathbb{Z} \to \tilde{C}^* \to C \to 1.$$

Suppose that $\tilde{C}^* \cap \mathbb{Z}$ is the subgroup $n\mathbb{Z}$ of $\mathbb{Z}$. Thus we have $\tilde{C}/\tilde{C}^* \simeq \mathbb{Z}/n\mathbb{Z}$.

In this case, $P_k : \tilde{G} \to \tilde{G}$ is dense if and only if k is coprime to n.

Indeed, as $P_k : \mathbb{Z}/n\mathbb{Z} \to \mathbb{Z}/n\mathbb{Z}$ is surjective if and only if k is coprime to n, the result follows from Theorem 1.4.

Case 2(b). C is disconnected, and $\pi_1(G) \simeq \mathbb{Z}$. Then there is a surjective group homomorphism $\tilde{C}/\tilde{C}^* \to C/C^*$. Let $D = \mathrm{Ker}(\tilde{C}/\tilde{C}^* \to C/C^*)$. As $\pi_1(G)$ is infinite cyclic, D is isomorphic to $\mathbb{Z}/n\mathbb{Z}$ for some n.

This happens for A_1I, C_nI ($n \geq 3$), D_nIII ($n \geq 4$ even).

Lemma 4.2. *In case 2(b), $P_k : \tilde{G} \to \tilde{G}$ is dense if and only if k is odd and coprime to n.*

Proof. First observe that, as C is disconnected, for even k, P_k is not dense.

In this case we get the following short exact sequence.

$$1 \to \mathbb{Z}/n\mathbb{Z} \to \tilde{C}/\tilde{C}^* \to C/C^* \to 1.$$

Let σ be any element in $\mathbb{Z}/n\mathbb{Z}$. If $\pi : \tilde{C}/\tilde{C}^* \to C/C^*$, then $\pi(\sigma) = 0$. Suppose that $\sigma = t^k$ for some $t \in \tilde{C}/\tilde{C}^*$. Then we have $0 = \pi(\sigma) = \pi(t^k) = \pi(t)^k$. As for odd integer, P_k is identity on C/C^*, we conclude that $t \in \mathbb{Z}/n\mathbb{Z}$. Therefore, if k is not coprime to n, P_k is not surjective on $\tilde{C}/\tilde{C}^*$. On the other hand, if k is odd and coprime to n, then both

$P_k : \mathbb{Z}/n\mathbb{Z} \to \mathbb{Z}/n\mathbb{Z}$ and $P_k : C/C^* \to C/C^*$ are surjective and the result follows from the short exact sequence. □

Case 3. C is not connected and $\pi_1(G) \simeq \mathbb{Z}/6\mathbb{Z}$. This happens for $E_6^2 II$.

Lemma 4.3. *$P_k : \tilde{G} \to \tilde{G}$ is dense if k is odd and coprime to 3 and for even k, $P_k : \tilde{G} \to \tilde{G}$ is not dense.*

Proof. We first note that, if k is odd and coprime to 3 (hence to 6), then P_k is surjective both on $\mathbb{Z}/6\mathbb{Z}$ and C. Hence P_k is surjective on $\tilde{C}$. Then the result follows by applying Theorem 1.1.

Since C is disconnected, P_2 can not be dense in $\tilde{G}$. □

Case 4. Let $\pi_1(G) \simeq \mathbb{Z}/4\mathbb{Z}$ or $\mathbb{Z}/8\mathbb{Z}$ and C is disconnected. This occurs for $D_n I(n \geq 4$ and n odd) and $D_n I(n \geq 4$ and n even) respectively. The following lemma is very straightforward and hence we omit the details.

Lemma 4.4. *In case 4, $P_k : \tilde{G} \to \tilde{G}$ is dense if and only if k is odd. Moreover, if G' be any cover of G, then $P_k : G' \to G'$ is dense if and only if k is odd.*

Case 5. G is split.

As usual, let $\tilde{G}$ be its universal cover. Let Z denote the discrete group $\pi_1(G)$. Suppose that there is a Cartan subgroup C of the form $A \times F$, where A is simply connected and F is a $\mathbb{Z}/2\mathbb{Z}$ vector space. Then we get the following short exact sequence,

$$1 \to Z \to \tilde{C} \to A \times F \to 1,$$

where $\tilde{C}$ is the corresponding Cartan subgroup of C. Let K be the kernel of the map $\tilde{C} \to A$. Then we have the following two short exact sequences

$$1 \to Z \to K \to F \to 1.$$

and

$$1 \to K \to \tilde{C} \to A \to 1.$$

Now, as A is simply connected, we get a section of the above short exact sequence, so that $\tilde{C} = A \ltimes K$. It turns out that $\tilde{C}$ is a direct product of A and K, as A is connected and K is discrete. Hence $P_k(\tilde{C}) = \tilde{C}$ if and only if $P_k(K) = K$. Furthermore, let $\sigma \in Z$. Let $\sigma = t^k$ for some $t \in K$. We denote π is the map from $K \to F$ in the above short exact sequence. Note that for even k, $P_k(\tilde{G})$ is not dense. For odd k, as $P_k : C \to C$ is identity,

$$0 = \pi(\sigma) = \pi(t^k) = P_k(\pi(t)) = \pi(t),$$

we get $t \in Z$. Thus we have $P_k(\tilde{G})$ is dense in $\tilde{G}$ if and only if $P_k(Z) = Z$.

Let D be a subgroup of Z and $G' = \tilde{G}/D$. Then by above procedure we can conclude that $P_k(G')$ is dense in G' if and only if $P_k(Z/D) = Z/D$.

Remark 4.5. In Case 3, the authors have not addressed the case when k is odd and divisible by 3, as they are not able to calculate the number of connected components of the Cartan subgroups involved. They have also left unaddressed the case when G is a non-split, non-compact simple Lie group and $\pi_1(G)$ contains an infinite cyclic subgroup, for the same reason.

5. Relation between square map and weak exponentiality

In this section we discuss the relation between the density of P_2 and weak exponentiality of Lie groups. Also, we prove Corollary 1.5.

Proof of Corollary 1.5: Let R be the solvable radical of G. Then G/R is a connected linear semisimple Lie group. As any connected semisimple Linear Lie group is precisely the Hausdorff connected component of some Zariski connected semisimple real algebraic group, Cartan subgroups of a connected linear Lie group are either connected or it has 2^m (for some positive integer m) number of disconnected components. Let C be any Cartan subgroup and C^* be its connected component. As $P_2(G)$ is dense in G, by Corollary 1.4, we get $P_2 : C/C^* \to C/C^*$ is surjective. This implies that $C = C^*$, and hence G is weakly exponential.

The converse follows from Corollary 1.3. □

Example 5.1. We give an example of a non-linearizable Lie group G which is not weakly exponential, but $P_2(G)$ is dense. Let us consider $G = \mathrm{PSL}(2,\mathbb{R})$ and $\tilde{G} = \widetilde{\mathrm{SL}(2,\mathbb{R})}$. We know that $Z(\tilde{G})$ is infinite cyclic. Fix a generator σ. Let Z to be the subgroup of $Z(\tilde{G})$ generated by 3σ and define $G' = \tilde{G}/Z$. There is a Cartan subgroup of G isomorphic to $\mathbb{R}$, and the corresponding Cartan subgroup for $\tilde{G}$ is isomorphic to $\mathbb{R} \times \mathbb{Z}$, where the factor $\mathbb{Z}$ is contributed by the center. Therefore, the corresponding Cartan subgroup of G' would be isomorphic to $\mathbb{R} \times \mathbb{Z}/3\mathbb{Z}$, which is not connected. By Theorem 1.1, $P_2(G')$ is dense in G', but it has a disconnected Cartan subgroup, and hence G' is not weakly exponential.

Remark 5.1. Let G be a connected Lie group, not necessarily linear, and let R be its radical. Suppose the adjoint semisimple group $\mathrm{Ad}(G/R)$ does not contain any simple factor whose fundamental group contains an infinite cyclic subgroup. It follows from the cases discussed in the previous section (§4) that if $P_2(G)$ is dense then G is weakly exponential.

6. Density on full rank subgroup

We recall that a Lie subgroup A of G is of *full rank* if its Lie algebra $\mathfrak{a}$ contains a Cartan subalgebra of $\mathfrak{g} = \mathrm{Lie}(G)$. In this case, every Cartan subalgebra of $\mathfrak{a}$ is a Cartan subalgebra of $\mathfrak{g}$ ([B, Proposition 3, §3, Ch. VII]).

Proposition 6.1. *Let G be a connected Lie group and let A be a connected subgroup of full rank. Let $k \geq 1$ be an odd integer. If $P_k(G)$ is dense in G, then $P_k(A)$ is dense in A.*

Proof. Note that since A is of full rank, $Z(G)^* \subset A$. By killing $Z(G)^*$, it is easy to see that $A_1 = A/Z(G)^*$ is a full rank subgroup of $G_1 = G/Z(G)^*$. Inductively, set $Z_i = Z(G_i)^*$ and $A_i = A_{i-1}/Z_{i-1}$. Then A_i is of full rank in G_i. Since the dimension of G is finite, Z_i is discrete for some i. It is enough to show that $P_k(A_i)$ is dense in A_i. For, if H is a connected Lie group and $Z \subset H$ is central connected, then $P_k(H)$ is dense in H if and only if $P_k(H/Z)$ is dense in H/Z. This is because, the projection $\pi : H \to H/Z$ is a submersion, hence an open map, and $\pi^{-1}(P_k(H/Z)) = P_k(H)$.

Thus we can assume that $Z(G)$ is discrete. Since $Z(G)$ is the fundamental group of linear Lie group $\mathrm{Ad}(G)$, it is a finitely generated abelian group.

We know that $\mathrm{Reg}(G) \cap A$ is dense in A (see proof of [N, I.6]). It is enough to prove that elements in $\mathrm{Reg}(G) \cap A$ admit k-th roots in A. Let $g \in \mathrm{Reg}(G) \cap A$. There is a unique Cartan subgroup C of G such that $g \in C$. The subgroup $C' := C \cap A$ contains the identity component C^*, which is also the identity component of the Cartan subgroup of A that contains g, with Lie algebra given by $N_{\mathfrak{a}}(Ad_g - I) = N_{\mathfrak{g}}(Ad_g - I)$. Therefore, C' is the disjoint union of cosets of C^*, hence open in C'. It is enough to show that $P_k(C') = C'$.

Writing $D = Z(G)$, let $\pi : G \to G/D$ be the projection. Now, by Theorem 1.1, $P_k(C) = C$. We get an exact sequence as follows, where E is the corresponding Cartan subgroup of G/D:

$$1 \to D \to C \xrightarrow{\pi} E \to 1.$$

This gives another short exact sequence

$$1 \to D/(D \cap C^*) \to C/C^* \xrightarrow{\pi} E/E^* \to 1.$$

However, E/E^* is an $\mathbb{Z}/2\mathbb{Z}$-vector space, so for odd $k \geq 0$, P_k induces identity on E/E^*. Therefore P_k is surjective on $D/(D \cap C^*)$. Indeed, let $x \in D/(D \cap C^*)$. Since $P_k(C) = C$, there is some $y \in C/C^*$ such that $x = y^k$. But $0 = \pi(x) = \pi(y)^k = \pi(y)$, so $y \in D/(D \cap C^*)$. This means, $D/(D \cap C^*)$ is a torsion abelian group with torsion coprime to k. However, C' fits into a short exact sequence

$$1 \to D' \to C' \to F \to 1,$$

where $F = \pi(C')$ is an open subgroup of E, and $D' = D \cap C'$. This gives rise to the short exact sequence as follows.

$$1 \to D'/(D' \cap C^*) \to C'/C^* \to F/F^* \to 1.$$

Here F/F^* is the group of connected component of $F \subset E$, which must be an $\mathbb{F}_2$-vector space, and $D'/(D' \cap C^*)$ is a subgroup of $D/D \cap C^*$, hence a torsion abelian group with torsion coprime to k. Therefore P_k is surjective on C'/C^*. □

Lemma 6.2. *Let G be a connected Lie group and let A be a connected full rank subgroup. Let C be any Cartan subgroup and C^* be its connected component. Suppose that C/C^* is finitely generated abelian. Let $k > 0$ be an integer. If $P_k : G \to G$ is dense then $P_k : A \to A$ is dense.*

Proof. In view of Proposition 6.1, it is enough to prove for integer $k = 2$. As $P_2(G)$ is dense in G, by Corollary 1.4, $P_2 : C/C^* \to C/C^*$ is surjective. By hypothesis C/C^* is a torsion abelian group and torsion is coprime to 2. We recall from the proof of Proposition 6.1 that, to prove $P_k : A \to A$ is dense, it only needs to show that $P_2 : A \cap C/C^* \to A \cap C/C^*$ is surjective. Note that $A \cap C/C^*$ is also torsion subgroup of C/C^* and torsion is coprime to 2 as well. This proves the lemma. □

Proposition 6.3. *Let G be a connected Lie group and let A be a connected full rank subgroup of G. Then $P_2 : G \to G$ is dense implies $P_2 : A \to A$ is dense.*

Proof. Arguing as before (see the proof of Proposition 6.1), we may assume that $Z(G)$ is discrete. Now, let $\pi : G \to \mathrm{Ad}(G)$ be the covering map and C be a Cartan subgroup of G. Let $C_{\mathrm{Ad}} = \pi(C)$ be the corresponding Cartan subgroup of $\mathrm{Ad}(G)$. Since $P_2(G)$ is dense in G, $P_2 : \mathrm{Ad}(G) \to \mathrm{Ad}(G)$ is dense and hence C_{Ad} is connected. Let C^* denote the identity component of C. We observe that the restriction map $\pi : C^* \to C_{\mathrm{Ad}}$ is surjective. Indeed, $\pi(C^*)$ is non empty open and closed in C_{Ad} (for details, see the proof of Lemma 2.3). Therefore we get a surjective homomorphism from $Z(G) \times C^* \to C_{\mathrm{Ad}}$. Hence C/C^* is finitely generated abelian group (as $Z(G)$ is finitely generated abelian group). Thus by Lemma 6.2, it follows that $P_2 : A \to A$ is dense. □

Proof of Theorem 1.6: The theorem follows from Proposition 6.1 and Proposition 6.3. □

Proof of Corollary 1.7: It is noted in Neeb ([N]) that, $Z_G(t_a)$ is a connected, full rank, reductive subgroup of G. Therefore forward implication of the statement follows from Theorem 1.6. Now for the converse, let C' be any Cartan subgroup of G. Let $\mathfrak{c}'$ be the Lie algebra associated to C'. Since $\mathfrak{g}$ is reductive, $\mathfrak{c}'$ is reductive in $\mathfrak{g}$ and its associated Cartan

subgroup is $C' = Z_G(\mathfrak{c}')$. We may assume that $\mathfrak{c}' = t'_a + \mathfrak{a}'$, where $\mathfrak{a}' \subseteq \mathfrak{a}$ and $t'_a \supseteq t_a$. Then it is immediate that $C' \subseteq Z_G(t_a)$ and by [N, Lemma I.5], C' is a Cartan subgroup of $Z_G(t_a)$. Now, by applying Theorem 1.1 on $Z_G(t_a)$, we conclude that $P_k(C') = C'$, as required. □

Acknowledgement. We thank Professor S. G. Dani for his useful comments, particularly Remark 3.1.

References

[B] Bourbaki, N. *Lie Groups and Lie Algebras*, Elements of Mathematics, Springer. Chapters 7-9.
[C1] Chatterjee, P. *On surjectivity of the power maps of solvable Lie groups*, J. Algebra 248 (2002), no. 2, 669 - 687.
[C2] Chatterjee, P. *On the power maps, orders and exponentiality of p-adic algebraic groups*, J. Reine Angew. Math., (2009), no. 629, 201 - 220.
[C3] Chatterjee, P. *Automorphism invariant Cartan subgroups and power maps of disconnected groups*, Math. Z. 269 (2011), no. 4, 221 - 233.
[DM] Dani, S.G. and Mandal, A. *On the surjectivity of the power maps of a class of solvable groups*, Preprint.
[F] Fell, J. *A Hausdorff topology for the closed subsets of a locally compact non Hausdorff space*,
[H] Hofmann, K.H. *A memo on the exponential function and regular points*, Arch. Math., (1992), no. 59, 24 - 37.
[H1] Hofmann, K.H. *Near-Cartan algebras and groups*, Seminar Sophus Lie, (1992), no. 2, 135 - 151.
[HL] K.H. Hofmann, J.D. Lawson, *Divisible subsemigroups of Lie groups*, J. London Math. Soc. 27 (1983), 427 - 437.
[HM] Hofmann, K.H., and A. Mukherjea, *On the density of the image of the exponential function*, Math. Ann., (1978), 263 -273.
[Mc] McCrudden, M. *On nth roots and infinitely divisible element*, Arch. Math., (1992), no. 59, 24 - 37.
[N] Neeb, K-H. *Weakly exponential Lie groups*, J. Algebra. 179 (1996), no. 15, 331-361.
[St] Steinberg, R. *On the power maps in algebraic groups*, Math. Res. Letters, (2003), no. 10, 621 - 624.
[W] Wüstner, M. *A generalization of the Jordan decomposition*. Forum Math. volume 15 (2003) 395-408.

Saurav Bhaumik
Department of Mathematics,
Indian Institute of Technology Bombay,
Powai, Mumbai 400076, India
saurav@math.iitb.ac.in

Arunava Mandal
Department of Mathematics,
Indian Institute of Technology Bombay,
Powai, Mumbai 400076, India
amandal@math.iitb.ac.in